Rhodora

JOURNAL OF THE
NEW ENGLAND BOTANICAL CLUB

NORTON H. NICKERSON, Editor-in-Chief JOAN Y. NICKERSON, Managing Editor

Associate Editors

DAVID S. BARRINGTON
A. LINN BOGLE
DAVID E. BOUFFORD
CHRISTOPHER S. CAMPBELL
WILLIAM D. COUNTRYMAN
GARRETT E. CROW

RICHARD A. FRALICK GERALD J. GASTONY C. BARRE HELLQUIST MICHAEL W. LEFOR ROBERT T. WILCE

VOLUME 94

1992

The New England Botanical Club, Inc.

Harvard University Herbaria, 22 Divinity Ave., Cambridge, Mass. 02138



INDEX TO VOLUME 94

New scientific names are in bold face

- A(risaema) dracontium, Arisaema triphyllum subsp. stewardsonii ×, hybrids in Massachusetts. A naturallyoccurring population of putative, 340–347
- Aconogonon (Polygonum sect. Aconogonon) in Alaska. Choosing the correct name for, 319–322
- Additions to the flora of Newfoundland. II. 383-386
- Alabama. A new species of Blephilia (Lamiaceae) from northern, 1-14
- Alaska. Choosing the correct name for Aconogonon (Polygonum sect. Aconogonon) in, 319–322
- Alpine flora of the northeastern United States. Contributions to the, 15–37
- Jesse M. Greenman Award 108, 399 NEBC Award for Support of Botanical Research 106, 397
- NEBC and New England Wildflower Society Symposium 99
- New England Botany Graduate Students Meeting 107
- Ronald L. Stuckey Herbarium Endowment Fund 107
- U.S. National Arboretum Herbarium 398
- Arethusa bulbosa, Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. Ecological aspects of, II. Partitioning the microhabitat. 374–380
- Arisaema triphyllum subsp. stewardsonii × A. dracontium hybrids in Massachusetts. A naturally-occurring population of putative, 340-347
- Armoracia lacustris (Brassicaceae) rediscovered in Ohio. 387–390
- Aster and Solidago. Chromosome number determinations in Fam. Compositae, Tribe Astereae. IV. Additional reports and comments on the

- cytogeography and status of some species of, 48-62
- Asteraceae of the Guianas, III: Vernonieae and the restoration of the genus Xiphochaeta. The, 348-361
- Barrington, David S. Climate and the disjunct distribution of Polystichum alfarii (Christ) Barr., comb. nov. in Mesoamerica. 327-339
- Berkshire County, Massachusetts. Natural plant communities of, 171–209
- Blephilia (Lamiaceae) from northern Alabama. A new species of, 1-14
- Blephilia subnuda Simmers & Kral, sp. nov. 2
- Boland, J. Todd and Peter J. Scott. Ecological aspects of Arethusa bulbosa, Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. II. Partitioning the microhabitat. 374–380

BOOK REVIEWS

- Phytogeography and Vegetation Ecology of Cuba. 103-105
- Plants of Northern British Columbia. 393–394
- Rare Plants of Prince Edward Island. The, 102
- Spores of the Pteridophyta: Surface, Wall Structure, and Diversity Based on Electron Microscope Studies. 100–101
- Bouchard, Andre see Hay, Stuart G.
- Brouillet, Luc see Hay, Stuart G. Burk, C. John see Sanders, Laurie L.
- Buttrick, Steven C. Habitat management: a decision making process. 258–286
- C.(hloris) dandeyana Adams. C. inflata Link and, Chloris barbata Sw. and C. elata Desvaux (Poaceae), the earlier names for, 135–140
- C.(hloris) elata Desvaux, Chloris bar-

bata Sw. and, (Poaceae), the earlier names for C. inflata Link and C. dandeyana Adams. 135-140

C.(hloris) inflata Link and C. dandeyana Adams. Chloris barbata Sw. and C. elata Desvaux (Poaceae), the earlier names for, 135–140

Caldwell, Fredricka A. and Garrett E. Crow. A floristic and vegetation analysis of a freshwater tidal marsh on the Merrimack River, West Newbury, Massachusetts. 63-97

Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. Ecological aspects of Arethusa bulbosa, II. Partitioning the microhabitat. 374–380

Canada. Geographical distribution and ecology of Long's Bulrush, Scirpus longii (Cyperaceae) in, 141-155

Cape Cod. An indigenous population of Clintonia borealis (Liliaceae) on, 98–99

Carex caryophyllea (Cyperaceae) in Massachusetts. Persistence of, 210– 211

Carolina Beach State Park (North Carolina). The flora of limesink depressions in, 156-166

Carpheporus corymbosus (Compositae). Comments on the phenology of, 323–325

Chambers, Kenton L. Choosing the correct name for Aconogonon (Polygonum sect. Aconogonon) in Alaska. 319–322

Chloris barbata Sw. and C. elata Desvaux (Poaceae), the earlier names for C. inflata Link and C. dandeyana Adams. 135–140

Chmielewski, Jerry G. see Semple, John C.

Choosing the correct name for Aconogonon (Polygonum sect. Aconogonon) in Alaska. 319-322

Chromosome number determinations in Fam. Compositae, Tribe Astereae. IV. Additional reports and comments on the cytogeography and status of some species of Aster and Solidago. 48-62

Chromosomes of Mexican Sedum VI. Section Sedastrum. 362-370

Climate and the disjunct distribution of Polystichum alfarii (Christ) Barr., comb. nov. in Mesoamerica, 327–339

Clintonia borealis (Liliaceae) on Cape Cod. An indigenous population of, 98-99

Comments on the phenology of Carpheporus corymbosus (Compositae). 323-325

Contributions to the alpine flora of the northeastern United States. 15-37

Corey, David T. Comments on the phenology of Carpheporus corymbosus (Compositae). 323–325

Cross, Shirley G. An indigenous population of Clintonia borealis (Liliaceae) on Cape Cod. 98–99

Crow, Garrett E. see Caldwell, Fredricka A.

Crow, Garrett E. see Weatherbee, Pamela B.

Cytogeography and status of some species of Aster and Solidago. Additional reports and comments on, Chromosome number determinations in Fam. Compositae Tribe Astereae. IV. 48–62

Distribution, disjunct, of Polystichum alfarii (Christ) Barr., comb. nov. in Mesoamerica. Climate and the, 327–339

Ecological aspects of Arethusa bulbosa, Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. II. Partitioning the microhabitat. 374–380

Endangered species. Scientific and policy considerations in restoration and reintroduction of, 287-315

Engelmannia peristenia (Raf.) Goodm. & Laws., comb. nov. 381

Etymology of Vaccinium L. On the, 371-373

- Falk, Donald A. and Peggy Olwell. Scientific and policy considerations in restoration and reintroduction of endangered species. 287–315
- Flora of limesink depressions in Carolina Beach State Park (North Carolina). The, 156-166
- Floral variation and taxonomy of Limnobium L. C. Richard (Hydrocharitaceae). 111-134
- Floristic and vegetation analysis of a freshwater tidal marsh on the Merrimack River, West Newbury, Massachusetts. A, 63-97
- Freshwater tidal marsh on the Merrimack River, West Newbury, Massachusetts. A floristic and vegetation analysis of a. 63–97
- Gandhi, Kancheepuram N. see Kartesz, John T.
- Geographical distribution and ecology of Long's Bulrush, Scirpus longii (Cyperaceae) in Canada. 141–155
- Goodman, George J. and Cheryl A. Lawson. Two new combinations and a name change from the Long Expedition of 1820. 381–382
- Guianas, Asteraceae of the, III: Vernonieae and restoration of the genus Xiphochaeta. The, 348-361
- Habitat management: a decision-making process. 258-286
- Hay, Stuart G., Andre Bouchard, Luc Brouillet and Martin Jean. Additions to the flora of Newfoundland. II. 383-386
- Herndon, Alan. Nomenclatural notes on North American Hypoxis (Hypoxidaceae). 43-47
- Hill, Nicholas M. and Mats E. Johansson. Geographical distribution and ecology of Long's Bulrush, Scirpus longii (Cyperaceae) in Canada. 141–155
- Holsinger, Kent E. Setting priorities for regional plant conservation programs. 243-257

- Hunt, David M. and Robert E. Zaremba. The northeastward spread of Microstegium vimineum (Poaceae) into New York and adjacent states. 167–170
- Hymenophyllum wilsonii Hooker (Hymenophyllaceae) in the Iberian peninsula. 316-318
- Hypoxis (Hypoxidaceae). Nomenclatural notes on North American, 43-47
- Iberian peninsula. Hymenophyllum wilsonii Hooker (Hymenophyllaceae) in the. 316-319
- IN MEMORIAM
- William H. Drury 395 Herman R. Sweet 396
- Indigenous population of Clintonia borealis (Liliaceae) on Cape Cod. An, 98–99
- Jean, Martin see Hay, Stuart G.
- Johansson, Mats E. see Hill, Nicholas M.
- Kartesz, John T. and Kancheepuram N. Gandhi. Chloris barbata Sw. and C. elata Desvaux (Poaceae), the earlier names for C. inflata Link and C. dandevana Adams. 135–140
- Keenan, Philip E. A new form of Triphora trianthophora (Swartz) Rydberg, and part 3 of observations of the ecology of Triphora trianthophora (Orchidaceae) in New Hampshire. 38-42
- Kral, Robert see Simmers, Richard W.
- Lawson, Cheryl A. see Goodman, George J.
- Limesink depressions in Carolina Beach State Park (North Carolina). The flora of. 156-166
- Limnobium L. C. Richard (Hydrocharitaceae). Floral variation and taxonomy of, 111-134
- Limnobium spongia (Bosc.) Steudel, subsp. laevigatum (Humboldt &

Bonplant ex Willdenow) Lowden, comb. nov. 129

Long Expedition of 1820. Two new combinations and a name change from the, 381-382

Long's Bulrush, Scirpus longii (Cyperaceae) in Canada. Geographical distribution and ecology of. 141–155

Lowden, Richard M. Floral variation and taxonomy of Limnobium L. C. Richard (Hydrocharitaceae). 111– 134

Marsh, freshwater tidal, on the Merrimack River, West Newbury, Massachusetts. A floristic and vegetation analysis of a, 63-97

Massachusetts. A floristic and vegetation analysis of a freshwater tidal marsh on the Merrimack River, West Newbury, 63–97

Massachusetts. A naturally-occurring population of putative Arisaema triphyllum subsp. stewardsonii × A. dracontium hybrids in, 340-347

Massachusetts. Natural plant communities of Berkshire County, 171–209 Massachusetts. Persistence of Carex caryophyllea in, 210–211

Massachusetts. Utricularia inflata Walter (Lentibulariaceae) in, 391-392

McCormac, James S. Armoracia lacustris (Brassicaceae) rediscovered in Ohio. 387–390

Merrimack River, West Newbury, Massachusetts. A floristic and vegetation analysis of a freshwater tidal marsh on the. 63-97

Mesoamerica. Climate and the disjunct distribution of Polystichum alfarii (Christ) Barr., comb. nov. in, 327– 339

Mexican Sedum VI, Chromosomes of, Section Sedastrum. 362–370

Microhabitat. Ecological aspects of Arethusa bulbosa, Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. II. Partitioning the, 374–380

Microstegium vimineum (Poaceae) into New York and adjacent states. The northeastward spread of, 167–170

Natural plant communities of Berkshire County, Massachusetts. 171–209

Naturally-occurring population of putative Arisaema triphyllum subsp. stewardsonii × A. dracontium hybrids in Massachusetts. A, 340–347 NEW ENGLAND NOTES

An indigenous population of Clintonia borealis (Liliaceae) on Cape Cod. 98–99

Persistence of Carex caryophyllea (Cyperaceae) in Massachusetts. 210-211

Utricularia inflata Walter (Lentibulariaceae) in Massachusetts. 391– 392

New Hampshire. A new form of Triphora trianthophora (Swartz) Rydberg, and part 3 of observations on the ecology of Triphora trianthophora (Orchidaceae) in, 38–42

New species of Blephilia (Lamiaceae) from northern Alabama. A, 1-14

New York and adjacent states. The northeastward spread of Microstegium vimineum (Poaceae) into, 167– 170

Newfoundland. Additions to the flora of, II. 383-386

Newfoundland. Ecological aspects of Arethusa bulbosa, Calopogon tuberosus and Pogonia ophioglossoides (Orchidaceae) in eastern, II. Partitioning the microhabitat. 374–380

Nomenclatural notes on North American Hypoxis (Hypoxidaceae). 43-47

North American Hypoxis (Hypoxidaceae). Nomenclatural notes on, 43-47

North Carolina. The flora of limesink depressions in Carolina Beach State Park, 156-166

Northeastward spread of Microstegium

vimineum (Poaceae) into New York and adjacent states. The, 167-170

Ohio. Armoracia lacustris (Brassicaceae) rediscovered in, 387–390 Olwell, Peggy see Falk, Donald A. Ortiz, S. see Soñora, F. X.

Persistence of Carex caryophyllea (Cyperaceae) in Massachusetts. 210– 211

Phenology of Carpheporus corymbosus (Compositae). Comments on the, 323–325

Plant conservation programs. Setting priorities for regional. 243-257

Pogonia ophioglossoides (Orchidaceae) in eastern Newfoundland. Ecological aspects of Arethusa bulbosa, Calopogon tuberosus and, II. Partitioning the microhabitat. 000–000

polystichum alfarii (Christ) Barr., comb. nov. 328

Polystichum alfarii (Christ) Barr., comb. nov. in Mesoamerica. Climate and the disjunct distribution of, 327–339

Rare species protection. Taxonomic issues in, 218–242

Restoration and reintroduction of endangered species. Scientific and policy considerations in, 287-315

Reviewers, List of, for 1991–1992.

Robinson, Harold. Asteraceae of the Guianas, III: Vernonieae and restoration of the genus Xiphochaeta. 348–361

Rodriguez-Oubiña, J. see Soñora, F. X.

Sanders, Laurie L. and C. John Burk. A naturally-occurring population of putative Arisaema triphyllum subsp. stewardsonii × A. dracontium hybrids in Massachusetts. 340-347

Scientific and policy considerations in restoration and reintroduction of endangered species. 287–315

Scirpus longii (Cyperaceae) in Canada. Geographical distribution and ecology of Long's Bulrush, 141-155

Scott, Peter J. see Boland, J. Todd Scutellaria parvula Michx. var. mis-

Scutellaria parvula Michx. var. missouriensis (Torr.) Goodm. & Laws., comb. nov. 381

Sedum, Chromosomes of Mexican, VI. Section Sedastrum. 362–370

Semple, John C., Jerry G. Climielewski and Chunsheng Xiang. Chromosome number determinations in Fam. Compositae, Tribe Astereae. IV. Additional reports and comments on the cytogeography and status of some species of Aster and Solidago. 48–62

Setting priorities for regional plant conservation programs. 243–257

Simmers, Richard W. and Robert Kral. A new species of Blephilia (Lamiaceae) from northern Alabama. 1–14

Siren, David J. and Karen R. Warr. The flora of limesink depressions in Carolina Beach State Park (North Carolina). 156–166

Solidago, Aster and, Chromosome number determinations in Fam. Compositae, Tribe Astereae. IV. Additional reports and comments on the cytogeography and status of some species of, 48-62

Soñora, F. X., S. Ortiz and J. Rodriguez-Oubiña. Hymenophyllum wilsonii Hooker (Hymenophyllaceae) in the Iberian peninsula. 316–319

Sorrie, Bruce A. Utricularia inflata Walter (Lentibulariaceae) in Massachusetts. 391-392

Standley, Lisa A. Persistence of Carex caryophyllea (Cyperaceae) in Massachusetts. 210-211

Standley, Lisa A. Taxonomic issues in rare species protection. 218–242

SYMPOSIUM-March, 1992.

New England Plant Conservation: The Scientific Basis for Effective Action. 213-315

Foreword. Leslie J. Mehrhoff, President NEBC and William E. Brum-

- back, Conservation Director NEWFS. 215-217
- Taxonomic issues in rare species protection. 218-242
- Triphora trianthophora (Swartz) Rydberg forma albidoflava Keenan forma nova. 39
- Triphora trianthophora (Swartz) Rydberg, A new form of, and part 3 of observations on the ecology of Triphora trianthophora (Orchidaceae) in New Hampshire. 38-42
- Two new combinations and a name change from the Long Expedition of 1820. 381-382
- Uhl, Charles H. Chromosomes of Mexican Sedum VI. Section Sedastrum. 362–370
- United States, northeastern. Contributions to the flora of the, 15-37
- Utricularia inflata Walter (Lentibulariaceae) in Massachusetts. 391–392

- Vaccinium L. On the etymology of, 371-373
- Vander Kloet, S. P. On the etymology of Vaccinium L. 371-373
- Vernonieae and restoration of the genus Xiphochaeta. The Asteraceae of the Guianas, III:, 348-361
- Warr, Karen R. see Siren, David J.
- Weatherbee, Pamela B. and Garrett E. Crow. Natural plant communities of Berkshire County, Massachusetts. 171-209
- Xiang, Chunsheng see Semple, John C. Xiphochaeta. The Asteraceae of the Guianas, III: Vernonieae and restoration of the genus, 348–361
- Zaremba, Robert E. see Hunt, David M.
- Zika, Peter F. Contributions to the flora of the northeastern United States. 15-37

